


# Divigel<sup>®</sup>


(estradiol hemihydrate)

## HOW TO APPLY DIVIGEL<sup>®</sup> GEL

- Always follow your healthcare professionals instructions on dosage and refer to the patient information leaflet for further information
- Store the gel at room temperature (below 25°C).




Apply the gel to a different side of your body each day. The Divigel<sup>®</sup> dose is applied once daily on the skin of the lower body or thighs.




Tear open the sachet.



Divigel<sup>®</sup> should be rubbed gently on dry and clean skin. The application surface should be 1-2 times the size of a hand.



Divigel<sup>®</sup> should not be applied on the breasts, on the face or irritated skin. Apply the gel to a different side of your lower body or thighs each day.



After application the gel should be allowed to dry for a few minutes and the application site should not be washed within 1 hour.



Contact of the gel with the eyes should be avoided. Hands should be washed after application. It should not be swallowed.

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Health Products Regulatory Authority (HPRA) website: [www.hpra.ie](http://www.hpra.ie) or email [medsafety@hpra.ie](mailto:medsafety@hpra.ie)

**ORION  
PHARMA**